

# Betadine Solution

<b>Drug Facts</b>	
<b>Active ingredient</b> Povidone-iodine, 10% (1% available iodine)	<b>Purpose</b> Antiseptic
<b>Uses</b> Patient pre-operative skin preparation <ul style="list-style-type: none"> <li>■ for preparation of the skin prior to surgery</li> <li>■ helps reduce bacteria that potentially can cause skin infection</li> </ul>	
<b>Warnings</b> For external use only <b>Do not use</b> ■ in the eyes <ul style="list-style-type: none"> <li>■ if you are allergic to povidone-iodine or any other ingredients in this preparation</li> </ul> <b>When using this product</b> <ul style="list-style-type: none"> <li>■ prolonged exposure to wet solution may cause irritation or, rarely, severe skin reactions</li> <li>■ in pre-operative prepping, avoid "pooling" beneath the patient</li> </ul> <b>Stop use and ask a doctor if</b> <ul style="list-style-type: none"> <li>■ irritation, sensitization, or allergic reaction occurs and lasts for 72 hours. These may be signs of a serious condition.</li> </ul> <b>Keep out of reach of children.</b> If swallowed, get medical help or contact a Poison Control Center right away.	
<b>Directions</b> <ul style="list-style-type: none"> <li>■ clean the operative site prior to surgery</li> <li>■ apply product and allow to dry</li> <li>■ may be covered with a bandage</li> </ul>	
<b>Other information</b> <ul style="list-style-type: none"> <li>■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)</li> <li>■ store in original container</li> </ul>	
<b>Inactive ingredients</b> pareth 25-9, purified water, sodium hydroxide	
<b>Questions?</b> 1-888-726-7535 (8am-5pm, EST, Mon.-Fri.).	

